varieties, competence & service



early firm cooking quality ware potato

- yellow flesh colour
- nice and good looking skin



Monique

early firm cooking nice skin quality ware potato

Variety characteristics				
Maturity	early		Tuber	
Cooking Type (EAPR)	AB		Tuber shape Eye depth Skin Flesh colour Raw discolouration	long oval - long shallow yellow - smooth yellow very low
Consumption quality	quality ware potato; very good taste; clear yellow flesh colour; no discolourati- on after cooking			
Yield	medium to high marketable yield; very good suitability for packing		Resistances PCN Potato Leafroll Virus Leaf blight Tuber blight Rhizoctonia Common scab	Ro 1 (9) high high-medium high high medium
Youth development	initially hesitant; later rapid			
Dormancy	high; excellent suitable for storage		Sensitivities Spraing Bruising Mechanical damage	low very low low
Growing recommendations				
Site requirements	medium to high demands to the soil and water supply, regular nutrient supply is recommended			
Planting	planting density	approx. 40.000 plants/ha (seed size 35/55 mm) => check QR Code for more details		
	planting depth	normal		
	seed preparation	normal compared with local standard		
	seed dressing	preventive control of <i>Rhizoctonia solani</i> is generally recommended		
Foliage	routine protective control of late blight is recommended			
Herbicides	no sensitivities known			
Fertilization further information				
Nitrogen	low-medium - compared with local standard			
Phosphorus	100 kg/ha (at planting, water soluble phosphate)			
Potassium	250 - 300 kg/ha			
Magnesium	80 kg/ha			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH

Printed on FSC certificated paper