
## varieties, competence & service



medium early, red skinned quality ware potato with firm texture

- nice oval tubers with remarkable intensive red skin
- fine grading, high marketable yield
- very good culinary quality



## Ramona

red skinned, medium early quality ware potato

**Ramona** - red skinned ware quality potato with remarkable external and internal quality

Variety characteristics				
Maturity	medium early		Tuber	
Cooking Type (EAPR)	A-B		Tuber shape Eye depth Skin Flesh colour	oval shallow red, smooth deep yellow
Consumption quality	quality ware potato with firm texture and pleasantly intensive flavour			
Yield	good tuber yield in a smooth grading		Resistances  Wart disease PCN Potato Virus y (PVY) Common scab	1 Ro 1(9), 3(9), 4(9), 5(9) high - very high medium - high
Youth development	rapid		Sensitivities	
Dormancy	high, good storability		Spraing Bruising Mechanical damage	low very low - low low - medium
Growing recommendations				
Site requirements	medium to high demands to soil quality and water supply			
Planting	planting density	~ 45,000 plants/ha (seed size 35/55 mm)		
	planting depth	normal		
	seed preparation	sprouting stimulation short before planting		
	seed dressing	preventive control Rhizoctonia solani is recommended; compatibility of particular active ingredients or products has to be checked		
Fertilization				
Nitrogen	local standard N-supply (160 kg/ha N-supply (incl. soil supplied N $_{\rm min}$ / incl. organic fertilizer); dose rates above 80 kg N/ha should be split on soils with risk of leaching			
Phosphorus	100 kg/ha (medium soil supply class; water soluble phosphate to planting)			
Potassium	250 - 300 kg/ha (up 150 kg/ha may be applied as potassium chloride)			
Magnesium	60 kg/ha			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.

## **EUROPLANT Pflanzenzucht GmbH**