
varieties, competence & service



very early, mainly firm cooking variety

- nice red skin colour with light yellow flesh
- good shape



Sanibel

very early mainly firm cooking red skin; variety with nice shape

Variety characteristics				
Maturity	very early	Tuber		
Cooking Type (EAPR)	В		Tuber shape	oval - round oval shallow red, smooth light yellow low
Consumption quality	light yellow flesh colo mainly firm cooking	our,	Eye depth Skin Flesh colour Raw discolouration	
Yield	high marketable yield; big tubers		Resistances Potato wart disease PCN Potato Leafroll Virus Leaf blight Tuber blight Rhizoctonia Common scab	D1 Ro 1(9) high high - medium high high
Youth development	rapid			
Dormancy	medium		Sensitivities	
			Spraing Bruising Mechanical damage	low - medium low low - medium
Growing recommendations				
Site requirements	medium demands to the soil and water supply			
Planting	planting density	approx. 40.000 plants/ha (seed size 35/55 mm) => check QR Code for more details		
	planting depth	normal		
	seed preparation	normal compared with local standard		
	seed dressing	preventive control of <i>Rhizoctonia solani</i> is generally recommended		
Foliage	routine protective control of late blight is recommended			
Herbicides	Sanibel has a slight sensitivity to herbicide Metribuzin; post-emergence application should be avoided			
Fertilization further information				
Nitrogen	medium - compared to local standard			
Phosphorus	100 kg/ha (at planting, water soluble phosphate)			
Potassium	200 - 250 kg/ha			
Magnesium	60 kg/ha			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH

Printed on FSC certificated paper